arXiv:1703.03590v1 [math.CV] 10 Mar 2017

First Order Differential Subordination for Functions with Positive Real Part

Om P. Ahuja, Sushil Kumar, and V. Ravichandran

Abstract. Sharp estimates on β are determined so that an analytic function p defined on the open unit disk in the complex plane normalized by $p(0) = 1$ is subordinate to some well known starlike functions with positive real part whenever $1 + \beta z p'(z)$, $1 + \beta z p'(z)/p(z)$, or $1 + \beta z p'(z)/p^2(z)$ is subordinate to $\sqrt{1+z}$. Our results provide sharp version of previously known results.

1. Introduction

Let $\mathcal{A}$ denote the class of analytic functions f on the disk $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ and normalized by the condition $f(0) = 0 = f'(0) - 1$. Let $\mathcal{S}$ be the subset of $\mathcal{A}$ of univalent functions. An analytic function f defined on $\mathbb{D}$ is subordinate to the analytic function g on $\mathbb{D}$ (or g is superordinate to f), if there exists an analytic function $w : \mathbb{D} \to \mathbb{D}$, with $w(0) = 0$, such that $f = g \circ w$. Furthermore, if g is univalent in $\mathbb{D}$, then $f \prec g$ is equivalent to $f(0) = g(0)$ and $f(\mathbb{D}) \subseteq g(\mathbb{D})$, see [**13**]. Let p be an analytic function on $\mathbb{D}$ normalized by $p(0) = 1$. Goluzin [**4**] discussed the first order differential subordination $zp'(z) \prec zq'(z)$ and proved that, whenever $zq'(z)$ is convex, the subordination $p(z) \prec q(z)$ holds and the function q is best dominant. After this basic result, many authors established several generalizations of first order differential subordination. The general theory of differential subordination is discussed in the monograph by Miller and Mocanu [**12**].

In 1989, Nunokawa *et al.* [**14**] proved that if subordination $1 + zp'(z) \prec 1 + z$ holds, then subordination $p(z) \prec 1 + z$ also holds. In 2007, Ali *et al.* [**2**] extended this result and determined the estimates on β for which the subordination $1 + \beta z p'(z)/p^j(z) \prec (1 + Dz)/(1 + Ez)$ $(j = 0, 1, 2)$ implies the subordination $p(z) \prec (1 + Az)/(1 + Bz)$, where $A, B, D, E \in [-1, 1]$. In 2013, Omar and Halim [**15**] determined the condition on β in terms of complex number D and real E with $-1 < E < 1$ and $|D| \leq 1$ such that $1+\beta z p'(z)/p^j(z) \prec (1 + Dz)/(1 + Ez)$ $(j = 0, 1, 2)$ implies $p(z) \prec \sqrt{1+z}$. Recently, Kumar and Ravichandran [**9**] determined some sufficient conditions for certain first order differential subordinations to imply that the corresponding analytic solution is subordinate to a rational, exponential, or sine function. For more details, see [**3, 19, 22, 24**]. The function $\sqrt{1+z}$ is associated with the class $\mathcal{S}^*_L$, introduced by Sokół and Stankiewicz [**23**]. This class consists of the function $f \in \mathcal{A}$ such that $w(z) := zf'(z)/f(z)$ lies in the region bounded by the right

2010 *Mathematics Subject Classification.* 30C45, 30C80.

Key words and phrases. Differential subordination; starlike function; functions with positive real part; Janowski function.

half of the lemniscate of Bernoulli given by $|w^2-1|<1$. The lemniscate of Bernoulli is a best known plane curve resembling the symbol ∞. It was named after James Bernoulli who considered it in elasticity theory in 1694. In geometry, the lemniscate is a plane curve defined by two given points F_1 and F_2, known as foci, at distance $2a$ from each other as the locus of points P so that $PF_1.PF_2=a^2$. The equation of lemniscate may be written as $(x^2+y^2)^2=2a^2(x^2-y^2)$. The lemniscate in the complex plane is the locus of $z=x+iy$ such that $|z^2-a^2|=a^2$.

For an analytic function $p(z)=1+c_1z+c_2z^2+\cdots$, we determine the sharp bound on β so that $p(z)\prec\mathcal{P}(z)$ where $\mathcal{P}(z)$ is a function with positive real part like $\sqrt{1+z}$, $(1+Az)/(1+Bz)$, e^z, $\varphi_0(z):=1+\frac{z}{k}\left((k+z)/(k-z)\right)$ $(k=\sqrt{2}+1)$, $\varphi_{sin}(z):=1+\sin z$, $\varphi_C(z):=1+\frac{4}{3}z+\frac{2}{3}z^2$ and $\varphi_{\leftmoon}(z):=z+\sqrt{1+z^2}$, whenever $1+\beta zp'(z)/p^j(z)\prec\sqrt{1+z}$, $(j=0,1,2)$. Many of our subordination results in this paper improve the corresponding non-sharp results obtained by earlier authors in [**1**, **6**, **11**]. Our results are sharp.

2. Main Results

In 1985, Padmanabhan and Parvatham [**16**] introduced a unified classes of starlike and convex functions using convolution with the function of the form $z/(1-z)^\alpha$, $\alpha\in\mathbb{R}$. Later, Shanmugam [**20**] considered the class $\mathcal{S}_g^*(h)$ of all $f\in\mathcal{A}$ satisfying $z(f*g)'/(f*g)\prec h$ where h is a convex function, g is a fixed function in $\mathcal{A}$. Denote by $\mathcal{S}^*(h)$ and $\mathcal{K}(h)$, the subclass $\mathcal{S}_g^*(h)$, when g is $z/(1-z)$ and $z/(1-z)^2$ respectively. In 1992, Ma and Minda [**10**] considered a weaker assumption that h is a function with positive real part whose range is symmetric with respect to real axis and starlike with respect to $h(0)=1$ with $h'(0)>0$ and proved distortion, growth, and covering theorems. The class $\mathcal{S}^*(h)$ generalizes many subclasses of $\mathcal{A}$, for example, $\mathcal{S}^*[A,B]:=\mathcal{S}^*((1+Az)/(1+Bz))$ $(-1\le B<A\le 1)$ [**5**], $\mathcal{S}_L^*:=\mathcal{S}^*(\sqrt{1+z})$ [**23**], $\mathcal{S}_e^*:=\mathcal{S}^*(e^z)$ [**11**], $\mathcal{S}_{sin}^*:=\mathcal{S}^*(\varphi_{sin}(z))$ [**7**], $\mathcal{S}_C^*:=\mathcal{S}^*(\varphi_C(z))$ [**21**], $\mathcal{S}_R^*:=\mathcal{S}^*(\varphi_0(z))$ [**8**], and $\mathcal{S}_{\leftmoon}^*:=\mathcal{S}^*(\varphi_{\leftmoon}(z))$ [**17**, **18**].

Several sufficient conditions for functions to belong to the above defined classes can be obtained as an application of the following subordination results involving the lemniscate of Bernoulli and other well known starlike functions with positive real part. Our first result gives a bound on β so that $1+\beta zp'(z)\prec\sqrt{1+z}$ implies that the function p is subordinate to several well-known starlike functions.

THEOREM 2.1. *Let the function p be analytic in $\mathbb{D}$, $p(0)=1$ and $1+\beta zp'(z)\prec\sqrt{1+z}$. Then the following subordination results hold:*

(a) *If $\beta\ge\frac{2(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))}{\sqrt{2}-1}\approx 1.09116$, then $p(z)\prec\sqrt{1+z}$.*
(b) *If $\beta\ge\frac{2(1-\log 2)}{3-2\sqrt{2}}\approx 3.57694$, then $p(z)\prec\varphi_0(z)$.*
(c) *If $\beta\ge\frac{2(1-\log 2)}{\sin(1)}\approx 0.729325$, then $p(z)\prec\varphi_{sin}(z)$.*
(d) *If $\beta\ge(2+\sqrt{2})(1-\log 2)\approx 1.044766$, then $p(z)\prec\varphi_{\leftmoon}(z)$.*
(e) *If $\beta\ge 3(1-\log 2)\approx 0.920558$, then $p(z)\prec\varphi_C(z)$.*
(f) *Let $-1<B<A<1$ and $B_0=\frac{2-\log 4-\sqrt{2}+\log(1+\sqrt{2})}{\sqrt{2}-\log(1+\sqrt{2}+1)}\approx 0.151764$. If either*
  (i) *$B<B_0$ and $\beta\ge\frac{2(1-B)(1-\log 2)}{A-B}\approx 0.613706\frac{1-B}{A-B}$ or*
  (ii) *$B>B_0$ and $\beta\ge\frac{2(1+B)(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))}{A-B}\approx 0.451974\frac{1+B}{A-B}$,*
  then $p(z)\prec(1+Az)/(1+Bz)$.

The bounds on β are sharp.

In proving our results, the following lemma will be needed.

LEMMA 2.2. *[**13**, Theorem 3.4h, p. 132] Let q be analytic in $\mathbb{D}$ and let ψ and ν be analytic in a domain U containing $q(\mathbb{D})$ with $\psi(w) \neq 0$ when $w \in q(\mathbb{D})$. Set $Q(z) := zq'(z)\psi(q(z))$ and $h(z) := \nu(q(z)) + Q(z)$. Suppose that (i) either h is convex, or Q is starlike univalent in $\mathbb{D}$ and (ii) $\operatorname{Re}(zh'(z)/Q(z)) > 0$ for $z \in \mathbb{D}$. If p is analytic in $\mathbb{D}$, with $p(0) = q(0)$, $p(\mathbb{D}) \subseteq U$ and*

$$\nu(p(z)) + zp'(z)\psi(p(z)) \prec \nu(q(z)) + zq'(z)\psi(q(z)),$$

then $p(z) \prec q(z)$, and q is best dominant.

PROOF OF THEOREM 2.1. The function $q_\beta : \overline{\mathbb{D}} \to \mathbb{C}$ defined by

$$q_\beta(z) = 1 + \frac{2}{\beta}(\sqrt{1+z} - \log(1+\sqrt{1+z}) + \log 2 - 1).$$

is analytic and is a solution of the differential equation $1 + \beta z q'_\beta(z) = \sqrt{1+z}$. Consider the functions $\nu(w) = 1$ and $\psi(w) = \beta$. The function $Q : \overline{\mathbb{D}} \to \mathbb{C}$ is defined by $Q(z) = zq'_\beta(z)\psi(q_\beta(z)) = \beta z q'_\beta(z)$. Since $\sqrt{1+z}-1$ is starlike function in $\mathbb{D}$, it follows that function Q is starlike. Also note that the function $h(z) = \nu(q_\beta(z)) + Q(z)$ satisfies $\operatorname{Re}(zh'(z)/Q(z)) > 0$ for $z \in \mathbb{D}$. Therefore, by making use of Lemma 2.2, it follows that $1+\beta zp'(z) \prec 1+\beta z q'_\beta(z)$ implies $p(z) \prec q_\beta(z)$. Each of the conclusion in (a)-(f) is $p(z) \prec \mathcal{P}(z)$ for appropriate $\mathcal{P}$ and this holds if the subordination $q_\beta(z) \prec \mathcal{P}(z)$ holds. If $q_\beta(z) \prec \mathcal{P}(z)$, then $\mathcal{P}(-1) < q_\beta(-1) < q_\beta(1) < \mathcal{P}(1)$. This gives a necessary condition for $p \prec \mathcal{P}$ to hold. Surprisingly, this necessary condition is also sufficient. This can be seen by looking at the graph of the respective functions.

(a) On taking $\mathcal{P}(z) = \sqrt{1+z}$, the inequalities $q_\beta(-1) \geq 0$ and $q_\beta(1) \leq \sqrt{2}$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = 2(1-\log 2)$ and $\beta_2 = 2(\sqrt{2} - 1 + \log 2 - \log(1+\sqrt{2}))/(\sqrt{2}-1)$ respectively. Therefore, the subordination $q_\beta(z) \prec \sqrt{1+z}$ holds only if $\beta \geq \max\{\beta_1,\ \beta_2\} = \beta_2$.

(b) Consider $\mathcal{P}(z) = \varphi_0(z)$. A simple calculation shows that the inequalities $q_\beta(-1) \geq \varphi_0(-1)$ and $q_\beta(1) \leq \varphi_0(1)$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = 2(1-\log 2)/(3-2\sqrt{2})$ and $\beta_2 = 2(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))$ respectively. Thus the subordination $q_\beta(z) \prec \varphi_0(z)$ holds only if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

(c) Consider $\mathcal{P}(z) = \varphi_{sin}(z)$. The inequalities $q_\beta(-1) \geq \varphi_{sin}(-1)$ and $q_\beta(1) \leq \varphi_{sin}(1)$ reduce to $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{2(1-\log 2)}{\sin(1)} \text{ and } \beta_2 = \frac{2(\sqrt{2} - 1 + \log 2 - \log(1+\sqrt{2}))}{\sin(1)}$$

respectively. The subordination $q_\beta(z) \prec \varphi_{Sin}(z)$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

(d) Consider $\mathcal{P}(z) = \varphi_{\mathbb{Q}}(z)$. The inequalities $q_\beta(-1) \geq \varphi_{\mathbb{Q}}(-1)$ and $q_\beta(1) \leq \varphi_{\mathbb{Q}}(1)$ give $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where $\beta_1 = (2+\sqrt{2})(1-\log 2)$ and $\beta_2 = \sqrt{2}(\sqrt{2} - 1 + \log 2 - \log(1+\sqrt{2}))$ respectively. The subordination $q_\beta(z) \prec \varphi_{\mathbb{Q}}(z)$ holds if $\beta \geq \max\{\beta_1, \beta_2\} = \beta_1$.

(e) Consider $\mathcal{P}(z) = \varphi_C(z)$. From the inequalities $\varphi_C(-1) \leq q_\beta(-1)$ and $q_\beta(1) \leq \varphi_C(1)$, we get $\beta \geq 3(1-\log 2)$ and $\beta \geq 2(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))$ respectively. Thus the subordination $q_\beta(z) \prec \varphi_C(z)$ holds if $\beta \geq \max\left\{3(1-\log 2),\ 2(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))\right\} = 3(1-\log 2)$.

(f) Consider $\mathcal{P}(z) = (1+Az)/(1+Bz)$. From the inequalities $q_\beta(-1) \geq (1-A)/(1-B)$ and $q_\beta(1) \leq (1+A)/(1+B)$, we note that $\beta \geq \beta_1$ and $\beta \geq \beta_2$, where

$$\beta_1 = \frac{2(1-B)(1-\log 2)}{A-B} \text{ and } \beta_2 = \frac{2(1+B)(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))}{A-B}$$

respectively. A simple calculation gives $\beta_1 - \beta_2 = 2(1-\log 2) + (1+B)(\log(1+\sqrt{2}) - \sqrt{2})$. We note that $\beta_1 - \beta_2 \geq 0$ if $B < B_0$ and $\beta_1 - \beta_2 \leq 0$ if $B > B_0$ where

$$B_0 = \frac{2 - \log 4 - \sqrt{2} + \log(1+\sqrt{2})}{\sqrt{2} - \log(1+\sqrt{2}+1)}.$$

The necessary subordination $p(z) \prec (1+Az)/(1+Bz)$ holds if $\beta \geq \max\{\beta_1, \beta_2\}$. ∎

The subordination results in part (a) and (f) in Theorem 2.1 were also investigated by the authors in [**1**, Lemma 2.1, p. 1019] and [**6**, Lemma 2.1, p. 3], but their results were non-sharp.

Next result gives a bound on β so that $1+\beta zp'(z)/p(z) \prec \sqrt{1+z}$ implies p is subordinate to some well-known starlike functions.

THEOREM 2.3. *Let the function p be analytic in $\mathbb{D}$, $p(0) = 1$ and $1 + \beta zp'(z)/p(z) \prec \sqrt{1+z}$. Then the following subordination results hold:*

(a) *If $\beta \geq \frac{2(\log 2 - 1)}{\log(2\sqrt{2}-2)} \approx 3.26047$, then $p(z) \prec \varphi_0(z)$.*
(b) *If $\beta \geq \frac{2\left(\sqrt{2}-1+\log(2)-\log\left(\sqrt{2}+1\right)\right)}{\log(1+\sin(1))} \approx 0.740256$, then $p(z) \prec \varphi_{sin}(z)$.*
(c) *If $\beta \geq \frac{2(\log 2-1)}{\log(\sqrt{2}-1)} \approx 0.696306$, then $p(z) \prec \varphi_{\mathbb{C}}(z)$.*
(d) *If $\beta \geq 2(1-\log 2) \approx 0.613706$, then $p(z) \prec e^z$.*
(e) *If $-1 < B < A < 1$ and $\beta \geq \max\{\beta_1, \beta_2\}$ where*

$$\beta_1 = \frac{2(1-\log 2)}{\log(1-B) - \log(1-A)} \text{ and } \beta_2 = \frac{2(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))}{\log(1+A)-\log(1+B)},$$

then $p(z) \prec (1+Az)/(1+Bz)$.

The bounds on β are best possible.

PROOF. The function $q_\beta : \overline{\mathbb{D}} \to \mathbb{C}$ defined by

$$q_\beta(z) = \exp\left(\frac{2}{\beta}(\sqrt{1+z} - \log(1+\sqrt{1+z}) + \log 2 - 1)\right)$$

is analytic and is a solution of the differential equation $1 + \beta zq'_\beta(z)/q_\beta(z) = \sqrt{1+z}$. Define the functions $\nu(w) = 1$ and $\psi(w) = \beta/w$. The function $Q : \overline{\mathbb{D}} \to \mathbb{C}$ defined by $Q(z) := zq'_\beta(z)\psi(q_\beta(z)) = \beta zq'_\beta(z)/q_\beta(z) = \sqrt{1+z} - 1$ is starlike in $\mathbb{D}$. The function $h(z) := \nu(q_\beta(z)) + Q(z) = 1 + Q(z)$ satisfies $\mathrm{Re}(zh'(z)/Q(z)) > 0$ for $z \in \mathbb{D}$. Therefore, by using Lemma 2.2, we see that the subordination

$$1 + \beta\frac{zp'(z)}{p(z)} \prec 1 + \beta\frac{zq'_\beta(z)}{q_\beta(z)}$$

implies $p(z) \prec q_\beta(z)$. As the similar lines of the proof of Theorem 2.1, the proofs of parts (a)-(e) are completed. ∎

The subordination in part (d) and (e) of Theorem 2.3 were earlier investigated in [**11**, Theorem 2.16(c), p. 10] and [**6**, Lemma 2.3, p. 5] where non-sharp results were obtained.

Next, we determine a bound on β so that $1+\beta zp'(z)/p^2(z) \prec \sqrt{1+z}$ implies p is subordinate to several well-known starlike functions.

THEOREM 2.4. *Let the function p be analytic in $\mathbb{D}$, $p(0)=1$ and $1+\beta zp'(z)/p^2(z) \prec \sqrt{1+z}$. Then the following subordination results hold for sharp bound of β:*

(a) *If $\beta \geq 4(1+\sqrt{2})(1-\log 2) \approx 2.96323$, then $p(z) \prec \varphi_0(z)$.*

(b) *If $\beta \geq \frac{2(1+\sin(1))(\sqrt{2}-\log(1+\sqrt{2})+\log 2-1)}{\sin(1)} \approx 0.989098$, then $p(z) \prec \varphi_{sin}(z)$.*

(c) *If $\beta \geq (2+\sqrt{2})(\sqrt{2}-\log(1+\sqrt{2})+\log 2-1) \approx 0.771568$, then $p(z) \prec \varphi_{\mathbb{Q}}(z)$.*

(d) *Let $-1<B<A<1$ and $A_0 = \frac{2-\log 4-\sqrt{2}+\log(1+\sqrt{2})}{\sqrt{2}-\log(1+\sqrt{2}+1)} \approx 0.151764$. If either*

(i) *$A>A_0$ and $\beta \geq \frac{2(1-A)(1-\log 2)}{A-B} \approx 0.613706\frac{1-A}{A-B}$ or*

(ii) *$A<A_0$ and $\beta \geq \frac{2(1+A)(\sqrt{2}-1+\log 2-\log(1+\sqrt{2}))}{A-B} \approx 0.451974\frac{1+A}{A-B}$,*

then $p(z) \prec (1+Az)/(1+Bz)$.

PROOF. The function $q_\beta : \overline{\mathbb{D}} \to \mathbb{C}$ defined by

$$q_\beta(z) = \left(1-\frac{2}{\beta}\left(\sqrt{1+z}-\log(1+\sqrt{1+z})+\log 2-1\right)\right)^{-1}$$

is clearly analytic and is a solution of the differential equation $1+\beta zq'_\beta(z)/q^2_\beta(z) = \sqrt{1+z}$. Define the functions $\nu(w)=1$ and $\psi(w)=\beta/w^2$. The function $Q : \overline{\mathbb{D}} \to \mathbb{C}$ defined by $Q(z) = zq'_\beta(z)\psi(q_\beta(z)) = \beta zq'_\beta(z)/q^2_\beta(z) = \sqrt{1+z}-1$ is starlike in $\mathbb{D}$, Q is starlike function. The function $h(z) := \nu(q_\beta(z)) + Q(z) = \nu(q_\beta(z)) + Q(z)$ satisfies the inequality $\mathrm{Re}(zh'(z)/Q(z)) > 0$ for $z \in \mathbb{D}$. Therefore, by using Lemma 2.2, we see that the subordination

$$1+\beta\frac{zp'(z)}{p^2(z)} \prec 1+\beta\frac{zq'_\beta(z)}{q^2_\beta(z)}$$

implies $p(z) \prec q_\beta(z)$. As the similar lines of the proof of Theorem 2.1, the proofs of parts (a)-(d) are obtained. ∎

The subordination in part (d) of Theorem 2.4 was earlier investigated in [**6**, Lemma 2.4, p. 6] where non-sharp result was obtained.

References

[1] R. M. Ali, N. E. Cho, V. Ravichandran and S. S. Kumar, Differential subordination for functions associated with the lemniscate of Bernoulli, Taiwanese J. Math. **16** (2012), no. 3, 1017–1026.

[2] R. M. Ali, V. Ravichandran and N. Seenivasagan, Sufficient conditions for Janowski starlikeness, Int. J. Math. Math. Sci. **2007**(2007), Art. ID 62925, 7 pp.

[3] N. E. Cho, H. J. Lee, J. H. Park and R. Srivastava, Some applications of the first-order differential subordinations, Filomat **30** (2016) no. 6, 1465–1474.

[4] G.M. Goluzin, On the majorization principle in function theory, Dokl. Akad. Nauk. SSSR, 42(1935), pp. 647–650.

[5] W. Janowski, Extremal problems for a family of functions with positive real part and for some related families, Ann. Polon. Math. **23** (1970/1971), 159–177.

[6] S. Sivaprasad Kumar, V. Kumar, V. Ravichandran and N. E. Cho, Sufficient conditions for starlike functions associated with the lemniscate of Bernoulli, J. Inequal. Appl. **2013**(2013), 176, 13 pp.

[7] S. Sivaprasad Kumar, V. Kumar and V. Ravichandran, Radius problem for sin-starlike functions, submitted.
[8] S. Kumar and V. Ravichandran, A subclass of starlike functions associated with a rational function, Southeast Asian Bull. Math. **40** (2016) no. 2, 199–212.
[9] S. Kumar and V. Ravichandran, Subordinations for functions with positive real part, preprint.
[10] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, 157–169, Conf. Proc. Lecture Notes Anal., I, Int. Press, Cambridge, MA.
[11] R. Mendiratta, S. Nagpal, and V. Ravichandran, On a subclass of strongly starlike functions associated with exponential function, Bull. Malays. Math. Sci. Soc. **38** (2015), no. 1, 365–386.
[12] S. S. Miller and P. T. Mocanu, On some classes of first-order differential subordinations, Michigan Math. J. **32** (1985), no. 2, 185–195.
[13] S.S. Miller and P. T. Mocanu, *Differential Subordinations*, Dekker, New York, 2000.
[14] M. Nunokawa, M. Obradović and S. Owa, One criterion for univalency, Proc. Amer. Math. Soc. **106** (1989), no. 4, 1035–1037.
[15] R. Omar and S. A. Halim, Differential subordination properties of Sokół-Stankiewicz starlike functions, Kyungpook Math. J. **53** (2013), no. 3, 459–465.
[16] K. S. Padmanabhan and R. Parvatham, Some applications of differential subordination, Bull. Austral. Math. Soc. **32** (1985), no. 3, 321–330.
[17] R. K. Raina and J. Sokół, On coefficient estimates for a certain class of starlike functions, Hacet. J. Math. Stat. **44** (2015), no. 6, 1427–1433.
[18] R. K. Raina and J. Sokół, Some properties related to a certain class of starlike functions, C. R. Math. Acad. Sci. Paris **353** (2015), no. 11, 973–978.
[19] V. Ravichandran and K. Sharma, Sufficient conditions for starlikeness, J. Korean Math. Soc. **52** (2015) no. 4, 727–749.
[20] T. N. Shanmugam, Convolution and differential subordination, Internat. J. Math. Math. Sci. **12** (1989), no. 2, 333–340.
[21] K. Sharma, N. K. Jain, and V. Ravichandran, Starlike functions associated with a cardioid, Afr. Mat. **27** (2016), no. 5-6, 923–939.
[22] K. Sharma and V. Ravichandran, Applications of subordination theory to starlike functions, Bull. Iranian Math. Soc. **42** (2016) no. 3, 761–777.
[23] J. Sokół and J. Stankiewicz, Radius of convexity of some subclasses of strongly starlike functions, Zeszyty Nauk. Politech. Rzeszowskiej Mat. No. 19 (1996), 101–105.
[24] N. Tuneski, T. Bulboacă, and B. Jolevska-Tunesk, Sharp results on linear combination of simple expressions of analytic functions, Hacet. J. Math. Stat. **45** (2016), no. 1, 121–128.

Department of Mathematics, Kent State University, Burton, USA
E-mail address: oahuja@kent.edu

Bharati Vidyapeeth's College of Engineering, Delhi–110063, India
E-mail address: sushilkumar16n@gmail.com

Department of Mathematics, University of Delhi, Delhi–110007, India
E-mail address: vravi@maths.du.ac.in; vravi68@gmail.com